

DESCRIPTION

STEALTH VIRUS DETECTION IN THE CHRONIC FATIGUE SYNDROME

Sub A1
00085271.052698

This application is a continuation-in-part of co-
pending United States patent application Serial No.
07/887,502, filed May 22, 1992 entitled "Stealth Virus
Detection in the Chronic Fatigue Syndrome" which is a
continuation-in-part application of United States patent
applications Serial No. 07/704,814, filed May 23, 1991,
and Serial No. 07/763,039, filed September 20, 1991
entitled "Spumavirus detection in the chronic fatigue
syndrome". These prior submissions, including any
drawings, are incorporated by reference in their entirety
herein.

1. Field of Invention

The present invention relates generally to methods
for detecting the presence of a virus. More particularly,
it relates to the detection of viruses by tissue culture
techniques.

2. Background of the Invention

A. Chronic Fatigue Syndrome

Palca, Science, 249:1240-1241 (1990) and Palca,
Science, 254:1726-1728 (1991) describe attempts to
identify a causative agent for chronic fatigue syndrome.

DeFreitas et al., Chemical Abstracts, 114: Abstract
No. 205331c (1991) describes retroviral sequences related
to human T lymphotropic virus type 2 in patients with
chronic fatigue immune dysfunction syndrome.

Gupta et al., Scandinavian Journal of Immunology,
33:319-327 (1991) describes a comprehensive immunological
analysis of chronic fatigue syndrome. The analysis of
cell mediated and antibody mediated immunity was performed
in 20 patients with chronic fatigue syndrome and 20 age
and sex matched healthy controls.

2

B. Culture Techniques

Werner, *Lancet*, II:258-259 (1979) describes the isolation of foamy viruses from patients with de Quervain Thyroiditis and the detection of a cytopathic effect.

- 5 Freshney, *Culture of Animal Cells: A Manual of Basic Technique*, pp. 127-136 (Alan R. Liss Inc., 1987) describes the maintenance of cell cultures and states that four factors indicate the need for replacement of culture medium: (1) a drop in pH; (2) high cell concentration;
10 (3) cell types such as transformed cells, continuous cell lines and some embryonic cells that deteriorate rapidly at high cell densities; and (4) morphological deterioration of the cell such as granularity around the nucleus, cytoplasmic vacuolation, and rounding up of the cells with
15 detachment from the substrate.

DiLuca et al., *Virology*, 175:199-210 (1990) describes the replication of viral and cellular DNA in human herpesvirus 6-infected cells and the use of medium RPMI 1640 containing 10% fetal calf serum.

- 20 Ablashi et al., *International Journal of Cancer*, 42:787-791 (1988) describes the use of human hematopoietic cell lines for the propagation of HBLV (human herpesvirus 6) in RPMI 1640 supplemented with 10% FBS and antibiotics.

- Rethwilm et al., *Nucleic Acids Research*, 18:733-738
25 (1990) describes an infectious molecular clone (pHSRV) of the human spumaretrovirus (HSRV). pHSRV derived virus produced foamy virus typical cytopathic effects in susceptible cultures.

09085271-052698

3. Summary of the Invention

The present invention provides a method for detecting a stealth virus by culturing a sample under conditions that induce a cytopathic effect. Several culture
5 conditions are sufficient to induce a cytopathic effect in a sample containing a stealth virus. These culture conditions include replacing the culture medium every 24 to 72 hours, adding 5% to 10% fetal calf serum to the culture medium, using serum free medium X Vivo-15, using
10 preculture centrifugation and adding viral enhancing medium to the culture.

In one aspect, a method of detecting a stealth virus is provided by culturing a sample under conditions in which any stealth virus in the sample is able to induce a
15 cytopathic effect.

The term "stealth virus" refers to a virus having all of the following characteristics: (a) the ability to induce a cytopathic effect in fibroblastic cultures including primary kidney cell cultures which is
20 characterized by the production of foamy appearing cells, including cell syncytia; (b) the ability to produce a toxin capable of suppressing viral growth; (c) the ability to grow in cells from a plurality of species; (d) the inability of viral infected cells to react in a typical
25 manner using typing antisera specific for cytomegalovirus, herpes simplex virus, varicella zoster virus, Epstein-Barr virus and human T cell lymphotropic virus (HTLV); (e) the inability of viral infected cells to hybridize in a typical manner with nucleic acids probes specific for
30 HTLV, cytomegalovirus, herpes simplex virus, human herpes virus-6, varicella zoster virus and Epstein-Barr viruses using stringent hybridization conditions; and (f) the inability to evoke an inflammatory response in tissues which it infects.

35 The term "cytopathic effect" (CPE) refers to the appearance of rounded, slightly enlarged, refractile cells

4

throughout the culture. In some cultures the CPE progresses to very prominent collections of tightly packed, enlarged, foamy-cell appearing cells, with clearly defined cell syncytia and evidence of considerable cell
5 destruction. Stages between the spindle shape of the normal fibroblasts and the rounded appearance of affected cells can be seen. Several inclusions, consistent with vacuoles, can be seen within the cytoplasm. As their numbers increase, affected cells form several tightly
10 adherent clumps with indistinct cell boundaries. The affected cells continue to proliferate and scatter away from the cell clumps.

In preferred embodiments the presence of the stealth virus is detected by (1) inoculating a permissive cell
15 line with a sample, and (2) detecting *in vitro* a CPE in the permissive cell line; the cell line is maintained in culture medium that is replaced every 24 to 72 hours; the cell line is an insect cell line e.g., the cell line is derived from a plurality of species and may even include
20 *Spodoptera fruiperdo* derived ovarian cell line Sf9; and the sample is from a patient suspected of having chronic fatigue syndrome (CFS) based upon other recognized criteria well known to physicians in the art.

The term "chronic fatigue syndrome" (CFS) refers to
25 an illness whose major characteristic is an unexplained fatigue lasting beyond 6 months which results in greater than 50% reduction in an individual's normal level of activity (Holmes et al., "Chronic fatigue syndrome: A working case definition," *Ann. Intern. Med.*, 108:387-389
30 (1988); Holmes, "Defining the chronic fatigue syndrome", *Rev. Inf. Dis.*, 13 (Suppl. I):S53-5 (1991); Shafan, "The chronic fatigue syndrome" *Am. J. Med.*, 90:731-738 (1991)). To establish a clinical diagnosis, the patients should show evidence of suffering at least eight of the following
35 minor symptoms: fever, sore throat, myalgia, muscle weakness (which may be exacerbated by exercise),

5

arthralgia, lymphadenopathy, sleep disturbance, headaches, acute or subacute onset, and neuropsychological symptoms. The neuropsychological symptoms include a difficulty in thinking, dysnomia, confusion, forgetfulness, irritability, depression, photophobia and transient visual scotomata.

A cell is "permissive" if a particular virus causes a productive infection in it. A productive infection is a viral infection of a cell that produces progeny with the vegetative or lytic cycle. A productive infection by a stealth virus is characterized by the appearance of a cytopathic effect. Using appropriate conditions, stealth viruses infect and propagate in many mammalian cells *in vitro*, especially cultures of fibroblastic cells including primary kidney cell cultures, and other cell lines of epidermal, mesodermal, neuroectodermal and lymphoid origin, e.g., glial cells, myoblasts, etc. The distinctive CPE is, however, readily observed in primary fibroblast cultures. At present, therefore, these are the preferred cell lines to isolate the virus from clinical and animal samples. Suitable cells include human MRC-5 lung fibroblasts, primary human foreskin fibroblasts MRHF and rhesus monkey kidney RMK cells. These well known cell lines are available commercially (e.g., from Whittaker Bioproducts, Walkersville, Maryland). It should be noted that not all cell lines are permissive for growth and/or for the expression of a cytopathic effect (CPE). Conversely, other cell lines may be more efficient in promoting the growth of stealth viruses than the fibroblast cell lines currently used. The tissue culture cell lines are maintained in the laboratory using methods well known in the art.

In a second aspect a method of detecting a stealth virus is provided by: (1) inoculating a permissive cell line with a sample; (2) neutralizing stealth virus-

09085271 053699

6

associated toxic activity; and (3) detecting *in vitro* a cytopathic effect in the permissive cell line.

The term "stealth virus-associated toxin" or "toxin" refers to the molecular entity (or entities) that mediates the toxic activity observed in stealth virus cultures *in vitro*, and which appears capable of suppressing stealth virus growth. Cultures with toxin are often more difficult to infect with other stealth viral isolates. The toxic activity is detectable in serum and cerebrospinal fluid from stealth virus infected patients and may mediate certain symptoms associated with the patient's illness.

In preferred embodiments the toxic activity is neutralized by frequently replacing the culture medium, using medium in which the production of the toxin is minimized, or by adding a neutralizing agent to the culture; the neutralizing agent is fetal calf serum or retinoic acid, an antibody specific for the toxin, or a chemical or virus derived agent capable of competing with or reversing the toxic activity; and the sample is derived from a human, animal, food, or other environmental substance or object.

The term "neutralized" refers to any amount of inhibition or decrease. Thus, by neutralizing the toxic activity, one actually increases the detection of stealth virus. The term "replacing" refers to removing old culture medium and transferring new culture medium into the sample being tested to a new culture medium.

Stealth viruses may be isolated from a sample of blood or other biological samples including surgical and fine needle aspiration tissue biopsies, post mortem organ biopsies, throat swabs and saliva, urine, cerebrospinal fluid (CSF), other body fluids, blood and blood products intended for transfusion or for *in vitro* uses, vaccines, foods, and from the environment.

09085271-052698

In a third aspect, a method for detecting a stealth-virus is provided by: (1) inoculating a permissive cell line with a sample of the virus in a culture; (2) adding serum free medium to the culture; and (3) detecting in vitro the presence of a CPE in the permissive cell line.

In preferred embodiments, the serum free medium is medium X Vivo-15 (BioWhittaker, Inc., Walkersville, Maryland). X Vivo-15 is a Iscove's modified Delbecco's medium with the addition of human albumin, human insulin and human transferrin. Iscove's modification of Delbecco's medium is intended to support high density cell growth and has additional glucose, salts and vitamins compared to Delbecco's medium (Iscove and Melchers, *J. Experimental Medicine*, 147:923 (1963)). Albumin, insulin and transferrin are provided as a replacement for the need for serum. Since these proteins are of human origin, X Vivo-15 is suitable for growing human cells intended to be injected into autologous recipients.

Of several serum free media tested, medium X Vivo-15 was clearly superior to medium 199 plus 7% FCS for supporting stealth viral growth. The CPE developed more rapidly and there was less of a general loss of vitality seen with the usual stealth viral cultures. Medium X Vivo-15 was also found to be superior to other serum free media, for example, Aims-V medium from GIBCO BRL, Gaithersburg, Maryland. Aims-V medium also uses human albumin, insulin and transferrin, but the basic salt component is F-10 medium (Ham, *Exp. Cell Res.*, 29:515 (1963)), rather than Iscove's. X Vivo-15 was also superior to medium 199 plus bovine albumin, insulin and transferrin, with or without FCS.

In a fourth aspect a method of detecting a stealth virus is provided by: (1) co-centrifuging a sample of said virus with a permissive cell line of indicator cells; (2) inoculating the cell mixture into culture vessels; and (3) detecting in vitro a CPE in the permissive cell line.

09085271-052698

Pre-culture centrifugation (PCC) of patients' lymphoid cells with indicator fibroblast cells refers to co-centrifugation of patients' or animals' lymphocytes with freshly harvested fibroblasts and replating the cells
5 back into the original tubes containing the fibroblasts.

In preferred embodiments the method involves adding cytomegalovirus (CMV) supernatant, or viral enhancing medium, to the culture and frequently refeeding the culture medium. Viral enhancing medium (VEM) is derived
10 from the supernatants from actively replicating viruses which are functionally related to the stealth virus. Filtered, boiled medium collected from a CMV positive MRC-5 cell line showing a well defined 2+ CPE was able to supplement the stealth viral growth enhancing activity of
15 medium X Vivo-15 used alone and was designated VEM for "viral enhancing medium". Titration of this medium showed that 20%-30% was adequate to provide significant growth enhancement. X Vivo-15 medium containing 20% each of both CMV and HHV-6 supernatants was more effective than medium
20 containing 20% or 30% CMV supernatant.

In a fifth aspect, a method of detecting a stealth virus is provided by: (1) inoculating a permissive cell line with a sample of said virus in a culture; (2) adding viral enhancing medium to the culture; and (3) detecting
25 *in vitro* a CPE in the permissive cell line.

In preferred embodiments the viral enhancing medium contains 30% boiled, filtered products derived from the supernatant of cultures of cytomegalovirus and 70% medium X Vivo-15; and the cell line is maintained in a culture
30 medium, that is frequently replaced.

In a sixth aspect culturing a virus is provided by: (a) cocentrifuging a sample of said virus with a permissive cell line of indicator cells; (b) inoculating the cell mixture into culture vessels; (c) adding viral
35 enhancing medium to the culture; and (d) detecting *in vitro* a CPE in the permissive cell line.

9

The PCC step in combination with the use of VEM will also improve the detection of the CPE associated with cytomegalovirus (CMV) and human herpesvirus 6 (HHV-6). The growth of CMV and HHV-6, however, are less dependent
5 on these modifications than that of stealth viruses. Furthermore, frequent refeeding of the cultures is not nearly as important for these viruses as it is for stealth viruses.

In preferred embodiments the virus is a stealth
10 virus, cytomegalovirus, or human herpesvirus-6.

The summary of the invention described in detail above is not intended to limit in any way the scope of the present invention which is defined in the appended claims.

4. Detailed Description of the Invention

15 Preferred embodiments of the present invention are described in detail below. However, the following description of the preferred embodiments is not intended to limit in any way the scope of the present invention, which is defined in the appended claims.

20 The present invention provides several culturing conditions that induce a cytopathic effect in a sample containing a stealth virus. These culturing conditions include using approximately 5%-10% fetal calf serum, refeeding the culture medium every 24 to 72 hours using
25 viral enhancing medium and using preculture centrifugation. These conditions are important for inducing a cytopathic effect. For example, the failure to replace the culture medium every 24 to 72 hours often prevents detection of the cytopathic effect. Another
30 example is provided by the fact that not all serum free medium induce a cytopathic effect. Indeed, the use of a basal medium such as minimal essential medium with 2% fetal calf serum and weekly refeeding of the cultures as is commonly practiced in most clinical virology
35 laboratories will not yield a cytopathic effect with

09085271 052698

10

primary isolates of a stealth virus. Thus, it can be seen that only a carefully selected specific set of culturing conditions are capable of inducing a cytopathic effect in a sample containing a stealth virus.

5 5. Utility

 The present invention provides methods for the detection of a stealth virus. A virus is an infective agent and the stealth virus is associated with disease. The disease may be chronic fatigue syndrome (CFS) or one of several other diseases. Therefore, my invention has broad application to any area in which it is important to detect a stealth virus. Such areas include medical, veterinary, and agricultural diagnostics and industrial and pharmaceutical biological quality control.

 Many patients with a stealth viral infection have a stealth virus associated disease. Therefore, the detection of the presence of a stealth virus may allow one to confirm the diagnosis of a particular disease. Thus, those patients may avoid needless treatment, including psychiatric treatment. Indeed, it is useful to detect the stealth virus in order to develop a therapy or treatment for patients with a stealth viral associated disease. For example, in order to develop a vaccine it is useful to first detect the virus.

 In contrast, many patients without a stealth viral infection do not have a stealth virus associated disease. Thus, the failure to detect a stealth virus may indicate a psychiatric rather than physiological problem. Therefore, those patients may seek proper treatment, such as psychiatric treatment, rather than pursuing treatment aimed at eliminating or reducing the effects of a stealth virus.

 Since the stealth virus is associated with disease, it will often times be useful to detect the virus either in order to eliminate or avoid it. For example, in some

09085371-052696

[illegible]

The present invention is also useful in a variety of other ways readily apparent to those skilled in the art.

This invention will be more fully understood with reference to the examples which follow. The following examples are intended to illustrate the invention, but not to limit its scope which is defined in the claims appended hereto. The following examples are presented to illustrate the advantages of the present invention and to assist one of ordinary skill in the art in making and using the same, but are not intended in any way to otherwise limit the scope of the disclosure or the protection granted by a letters patent hereon.

12

multiple ~~or~~ enlarged refractable cells forming small clumps with indistinct cell borders are clearly seen in the cultures. A +/- (plus/minus) or equivocal effect refers to a minimal change in the culture in which the rounded cells are either not enlarged or non-refractile (possibly dead). Less than 4% of cultures from CFS patients have been recorded as equivocal, either because the CPE has been restricted to only a small region of the culture, or because it has failed to recur on secondary passage. These cultures have been excluded from analysis.

A ++ (two plus) strong positive response is recorded when vacuoles are clearly identified within enlarged, rounded, refractile cells and/or when multiple foci of cell clumping and/or destruction are apparent which are ringed by cells described as having a positive effect. A +++ (three plus) very strong positive effect refers to extensive ++ cellular changes throughout much of the culture and/or the appearance of large refractile vacuolated, foamy syncytial cell formation. Typical CPE's are shown in Patent Cooperation Treaty publication number WO 92/20787.

A benefit of using multiple indicator cell lines, is that the CPE appearances in the different cell lines sometime complement each other. For example, cell syncytia are usually best observed in monkey kidney cells. The smaller rounded refractile cells are usually best observed in the human fibroblasts. In over 80% of positive cultures, the CPE is clearly observable in at least 2 of the 3 cell lines. As with any viral induced CPE, it is important to confirm transmission to fresh cultures. In recipient cultures, the stealth virus induced CPE generally develops more rapidly than in the primary culture with at least the same level of intensity.

Although the appearances of the CPE share some features in common with those associated with cytomegalovirus (CMV) and herpes simplex virus (HSV) infection,

09085371-053998

stealth virus cultures can be readily distinguished from cultures harboring these viruses. The CPE from HSV is much more rapid, occurring in a matter of days. The destruction is greater with large masses of dead cells.

5 The CPE from CMV tends to initially develop in smaller, looser clusters and evolves more slowly than observed with stealth viruses. The intracytoplasmic vacuolization and syncytia formation are far less prominent with CMV than with stealth viruses. The stealth virus infected cells
10 give more the appearance of "foamy cells" than does CMV. Indeed, with some stealth viral cultures, there appears to be a marked accumulation of lipid-like material attaching itself to the wall of the culture tube. Human CMV does not infect monkey derived cells.

15 7. Example 1: Culture Of Stealth Viruses - 7% FCS and Frequent Refeeding

Culture tubes containing human fibroblast (MRC-5), primary human foreskin fibroblast (MRHF) and rhesus monkey kidney (RMK) cells are each inoculated with a cellular
20 mixture of the buffy coat granulocytes and ficoll-hypaque separated lymphocytes derived from approximately the equivalent of one milliliter (ml) of heparinized blood. Typically, 5 ml of blood collected into a "green top" heparinized tube, are layered onto 3 mls of a ficoll-
25 hypaque lymphocyte separation medium. Following 20 minutes centrifugation at 1,500 rpm, the lymphocyte, which collect at the plasma:ficoll hypaque interface, and the "buffy coat", present on the top of the erythrocyte layer, are collected into approximately 1 ml. An effort is made
30 to minimize the number of erythrocytes in collecting all of the visible buffy coat.

The cell mixture is washed once in 1 ml of 199 medium containing 7% FCS and resuspended into 1 ml. Aliquots of 0.2 ml of the cells are added to culture tubes containing
35 1 ml of 199 medium plus FCS. The tubes are placed in an

09085271-052698

14

incubator at 37°C for 45-60 minutes. The are then rinsed to remove macroscopically visible erythrocytes and other non-adherent cells. Rinsing (washing) consists of emptying the fluid content of the culture tube by decanting or by aspiration; adding 2 mls of medium or phosphate buffer saline; rocking the tube for several seconds to suspend erythrocytes; decanting the tube again. This important step is performed 2-10 times or until there are no macroscopically visible erythrocytes.

Two mls of medium with 7% FCS, are added and the tubes returned to a 37°C incubator. Cultures are maintained in the incubator at 37°C, with refeeding (replacement of old medium with medium) at 24, 48 and 72 hours. The tubes are examined microscopically after the 24 hour refeeding and, if residual erythrocytes are present, the tubes are rinsed in a manner similar to that performed at the 45-60 min. step. The tubes are refed three times each week by replacing the old medium with 2 mls of fresh medium. This procedure is designed to reduce the accumulation of a toxic component in the culture medium which tends to suppress viral growth. The tubes are examined three times per week for evidence of CPE (CPE). CPE is generally recognizable between 2-3 weeks after culturing.

A lack of regular refeeding of the cultures can result in a tendency for the CPE to abort and not to progress. This effect is not seen with CMV infected cultures. In unfed stealth virus cultures and even with regular refeeding, one can observe a toxic effect on many of the remaining cells. Culture cells appear to lose a degree of vitality and become duller in appearance compared to control cultures. The fibroblasts can assume somewhat of a pavement appearance, instead of the elongated shape. Some of these changes can be mimicked using 5 nM of the polyether marine toxin okadaic acid (Cohen et al., "Okadaic acid: A new probe for the study

09085271 052698

15

0607

Although presumptive of stealth virus infection, the
15 CPE appearance may require additional confirmation for a
definitive conclusion of stealth virus infection.

The medium was transferred to new tubes which were placed in a beaker containing water. The water was heated to boiling for 20 minutes. After cooling, the medium was
30 filtered through a 0.45 micron Millipore filter. This material was diluted 30:70 into regular X Vivo-15 medium to constitute a "lot" of CMV derived VEM. Each lot of VEM is tested to confirm: i) that it does not contain any residual live CMV by adding the medium to MRC-5 cells; and
35 ii) that it can promote the development of the CPE induced

by the prototype stealth virus by comparing the growth of the stealth virus in RMK cells containing VEM with that of similarly inoculated cells containing X Vivo-15 medium without supplement.

5 VEM has been tested on ten additional stealth viral isolates and clearly enhanced the growth of all of them. Two of these isolates are known to share CMV related sequences with the prototype stealth virus. Another isolate (from patient G.P.) shares antigens with HHV-6,
10 rather than CMV, and is considered an HHV-6 related stealth virus. Another isolate appears to have an adenoviral sequence. The use of VEM also reduced the time for a discernible CPE using fresh blood from two newly cultured CFS patients. It enhances the intensity of the
15 CPE and reduces the tendency for weekly positive cultures to revert to near normal appearance. VEM has also worked well in the cultures from the tissues of cats inoculated with the prototype stealth virus from patient D.W. allowing for clearly positive culture results. VEM
20 obtained from HHV-6 (strain GS) infected fibroblasts was similarly tested for its ability to promote the growth of CMV and HHV-6 associated stealth viruses. It worked well with both viral types with a discernable advantage on the HHV-6 related stealth virus from patient G.P., compared to
25 the CMV related stealth virus from patient D.W.

As a specific example, cultivation of a prototype stealth virus isolated from a CFS patient (initials D.W.) can be greatly enhanced by the addition to the culture of VEM comprising a 30% concentration of boiled, filtered
30 supernatants from cytomegalovirus (CMV) infected cultures. This addition helps remedy a deficiency of viral growth enhancing components coded for by the immediate-early (I-E) and probably other CMV related genes which are not detectable in the stealth virus from this patient.

09085271.052698

The lymphocyte-fibroblast cell mixture was centrifuged at high speed for 3 minutes. The tightly-packed cell pellet was gently resuspended and transferred back to the tube from which the fibroblasts were originally obtained. Two mls of VEM (X Vivo-15 medium supplemented with 30% CMV supernatant) were added and the tubes are placed in an incubator at 37°C. The tubes were refed with VEM at 48 and 72 hours and thereafter 3 times per week.

[illegible]

18

Control cultures in which either lymphocytes from other individuals are used, or the fibroblasts were processed but with the exception of no added lymphocytes, were fed in parallel with the test cultures. Note, in this revised protocol, buffy coat granulocytes are no longer routinely used since the contaminating erythrocytes tended to clump about the fibroblasts during the centrifugation step and were difficult to remove in subsequent washing of the cultures. Granulocytes may be an important source of virus in some patients. If this proves to be so, leucocyte rich plasma will be obtained by dextran precipitation of the erythrocytes from anti-coagulated blood, or as an alternative, modifications of the ficoll-hypaque separation method can be used which will separate both lymphocytes and granulocytes away from the erythrocytes. For example by using PMN isolation medium from Robbins Scientific Corp., Sunnyvale Ca. CSF and tissue extracts can be used in place of the lymphocytes.

The cultures were observed for a CPE which characteristically consists of rounding and swelling of the cells, formation of cell clumps which tend to disperse, appearance of intracellular granules/vacuoles and an overall foamy cell appearance often with prominent accumulation of lipid-like material.

09085271-053698

Table 3

Examples of the Enhanced Recovery and More Intense Development of CPE by a Stealth Virus from a CFS Patient Using Pre-Culture Centrifugation (PCC) and Viral Enhancing Medium (VEM).*

Method of Culturing	Time to CPE**	Intensity of CPE
Patient 1 Medium 199 + 7% FCS PCC and VEM	45 days 12 days	1-2+ 3+
Patient 2 Medium 199 + 7% FCS PCC and VEM	28 days 16 days	1-2+ 3+

* Medium X Vivo-15 containing 30% supplement of boiled filtered supernatant from a CMV culture also grown in medium X Vivo-15.

** Results are in RMK cells. Enhanced growth was also seen in MRC-5 cells.

10. Growth of Stealth Virus in Insect Cell Line

The Spodoptera fruiperda derived ovarian cell line Sf9 that is used routinely for the propagation of recombinant insect baculovirus was obtained from PharMigen, San Diego. It was maintained at 27°C in Grace's medium with 10% fetal calf serum. The stealth viruses from patients D.W., G.P., K.E. and B.B. were passaged into the insect cell line using 0.1 ml of cell-free supernatant from an infected MRC-5 human fibroblast culture. CPE was clearly seen within two days and progressed over the next several days.

The infected cultures showed enlarged foamy cell syncytia. Virus infectious for MRC-5 and for insect cell cultures was recoverable from the insect cell cultures to a dilution of 10^{-3} ml. Electron microscopic examination of the insect cultures infected with the virus from patient D.W. showed abundant herpes-like viral particles. In control studies, neither cytomegalovirus, human herpes virus 6, varicella zoster virus or Epstein-Barr virus

T200X

0908571 052698

20

induced CPE in the insect cell line; nor was infectious virus recoverable from these cultures.

11. Deposit Of Microorganisms

The stealth virus isolated from patient D.W. (virus-X
5 infected MRC-5 cells) was deposited with the American Type
Culture Collection (ATCC) - 12301 Parklawn Drive,
Rockville, Maryland 20852, under the provisions of the
Budapest Treaty on the International Recognition of the
Deposit of Microorganisms for the Purposes of Patent
10 Procedures on 9-17-91, and were assigned accession no. VR-
2343.

The present invention is not to be limited in scope
by the microorganisms deposited or the specific embodi-
ments described herein since such embodiments are intended
15 as but single illustrations of one aspect of the invention
and any microorganisms which are functionally equivalent
are within the scope of this invention. Indeed, various
modifications of the invention in addition to those shown
and described herein will become apparent to those skilled
20 in the art from the foregoing description and accompanying
drawings. Such modifications are intended to fall within
the scope of the appended claims.

Various references are cited herein, the disclosures
of which are incorporated by reference herein in their
25 entirety.

09085271.053698